



GREEN COW

feeding Instructions

WWW.SOFTAGRICULTURE.COM.AU

DIRECTIONS FOR USE

- This product is designed for the feeding of beef and dairy cattle.
- Offer according to the pasture quality, body weight and conditions of the animal.
- Introduce the ration gradually (over 3 - 4 weeks) to allow the rumen to adjust to the change in feed and to reduce the risk of ruminal acidosis.
- Feeders must not be allowed to be empty for a period of time. If feeders are empty for too long, hungry cattle may gorge themselves when feeders are re-filled, and will be at risk of acidosis.
- This product is not a complete feed. It must be fed in conjunction with quality hay and/or grazing. Roughage should make up no less than 20% of the animal's total dry matter intake.
- Fresh water must be provided at all times.

RECOMMENDED FEEDING RATE

Green Cow Weaner Pellet: 1 - 2 kg per head per day

Green Cow Breeder Pellet: 4 - 6 kg per head per day

Green Cow Dry Lick: 250 - 500 grams per head per day

TRANSPORTATION & STORAGE

Our products must be covered and protected from the rain at all times during transport. Shed storage is essential. Upon delivery, please remove plastic wrapping and covering from pallets, and leave bulk bags open for 24 hours to allow product to breathe. Stockfeed should always be protected against contamination from pests and rodents. Please note that failure to follow these guidelines may affect the quality of the product.

WEIGHT GAIN

Daily weight gain can be difficult to predict as it can vary considerably depending on the quantity and quality of pastures and hay available.